

A Pest in Paradise: Two-Line Spittle Bug Attacks Hawaii Rangelands
For Rangelands Partnership Newsletter, October 2, 2018
By Mark S. Thorne, Ph.D.
Range and Livestock Extension Specialist
University of Hawaii-Manoa
Cooperative Extension

Valued at more than \$68 million annually, livestock ranching is a highly productive and extremely important industry on all major islands in the State of Hawaii. Now however, the Hawaiian livestock industry faces severe threats from exotic pests such as the newly discovered two-lined spittlebug (TLSB), *Prosopia bicincta*. It is not known when the TLSB was introduced into Hawaii however, they were first detected in the Kailua-Kona area on Hawai'i Island in September of 2016 where it had caused severe damage to nearly 2,000 acres of rangeland. The pest has since rapidly expanding its range and is now causing similar damage to an estimated 45,000 acres. In highly infested areas there has been nearly 100% die back of the dominant and key range grasses. The loss of these grasses provides entry for the establishment of many undesirable and invasive plants. Currently TLSB appears to be isolated to the North and South Kona areas, but because of its seeming preference for key livestock forage grasses it has the potential to spread throughout the islands and irreparably harm large areas of valuable livestock grazing lands. There are currently no integrated pest management (IPM) protocols, nor are there any known natural enemies of TLSB present in Hawaii. Additionally, the potential geographic spread of the infestation is currently unknown but is potentially extensive, and very little is understood about how this new pest will perform in the Hawaiian environment. Consequently, TLSB is poised to become a major economic threat to the pasture-based livestock industry in Hawaii unless steps are taken to control the pest.

The native range of the Two-lined spittlebug extends from southeastern United States to Central America where it is a major pest of warm-season grasses. Adults resemble leafhoppers with two red stripes traversing the forewings held over the back of the body. The nymphs reside in "spittle masses" that protects them from desiccation and predators. The masses are found at the base of grasses near the soil line. Both nymphs and adults feed on the grass using a needle-like mouth part to penetrate the dermis and suck out the plant fluids. The adults may also inject the plants with *amylase*, an enzyme that breaks down starches. This feeding leads to weakened and stressed grasses that turn yellow as leaf material dies. Severe infestations can kill the grass.

To combat this pest a taskforce comprised of extension specialists and agents, graduate students and research technicians from the University of Hawaii and personnel from the Hawaii Dept. of Agriculture (HDOA) Plant Pest Control Branch was formed in the fall of 2017. Initial funding was provided by the HDOA, Dept. of Natural Resources, and Hawaii County. The goal of the taskforce is to restrict the further spread of TLSB and to prohibit its establishment on the other non-infested Hawaiian Islands through rancher outreach and education; detection and surveillance; development of Integrated Pest Management (IPM) protocols; biological control agent exploration; and research into the biology and ecology of TLSB in Hawaii.



Figure: Hawaii rangeland severely damaged by Two-line Spittlebug (*Prosapia bicincta*) resulting in subsequent infestation by fireweed (*Senecio madagascariensis*; green plants) and Pamakani (*Eupatorium adenophorum*; brown bush with white flowers).



Figure: Adult Two-line Spittlebug (*Prosapia bicincta*). Adults feed on the grasses using a needle-like mouth part to suck out the fluids from the plant.



Figure: Spittle masses of Two-line Spittlebug (*Prosapia bicincta*) nymphs along exposed roots of grasses. The nymphs feed on the juices from the plant for energy and to produce the spittle mass that protects them from desiccation and predators.