

**PROCEEDINGS OF THE AUSTRALIAN RANGELAND SOCIETY
BIENNIAL CONFERENCE**

Official publication of The Australian Rangeland Society

Copyright and Photocopying

© The Australian Rangeland Society 2012. All rights reserved.

For non-personal use, no part of this item may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior permission of the Australian Rangeland Society and of the author (or the organisation they work or have worked for). Permission of the Australian Rangeland Society for photocopying of articles for non-personal use may be obtained from the Secretary who can be contacted at the email address, rangelands.exec@gmail.com

For personal use, temporary copies necessary to browse this site on screen may be made and a single copy of an article may be downloaded or printed for research or personal use, but no changes are to be made to any of the material. This copyright notice is not to be removed from the front of the article.

All efforts have been made by the Australian Rangeland Society to contact the authors. If you believe your copyright has been breached please notify us immediately and we will remove the offending material from our website.

Form of Reference

The reference for this article should be in this general form;
Author family name, initials (year). Title. *In*: Proceedings of the nth Australian Rangeland Society Biennial Conference. Pages. (Australian Rangeland Society: Australia).

For example:

Anderson, L., van Klinken, R. D., and Shepherd, D. (2008). Aerially surveying Mesquite (*Prosopis* spp.) in the Pilbara. *In*: 'A Climate of Change in the Rangelands. Proceedings of the 15th Australian Rangeland Society Biennial Conference'. (Ed. D. Orr) 4 pages. (Australian Rangeland Society: Australia).

Disclaimer

The Australian Rangeland Society and Editors cannot be held responsible for errors or any consequences arising from the use of information obtained in this article or in the Proceedings of the Australian Rangeland Society Biennial Conferences. The views and opinions expressed do not necessarily reflect those of the Australian Rangeland Society and Editors, neither does the publication of advertisements constitute any endorsement by the Australian Rangeland Society and Editors of the products advertised.



The Australian Rangeland Society

IDENTIFYING STATE AND TRANSITION MODELS IN THE MITCHELL GRASSLANDS

David G. Phelps¹, Ockert J.H. Bosch² and Kirsten N. Bates¹

¹ Department of Primary Industries, PO Box 519 Longreach 4730 Qld Australia

² Landcare Research NZ LTD, PO Box 69 Lincoln 8152 Canterbury New Zealand

INTRODUCTION

The Mitchell grasslands are the most extensive and productive native pastures of semi-arid western Queensland. The grasslands are dominated by the long-lived perennial *Astrebula* spp. (Mitchell grasses). Rainfall is highly variable, leading to fluctuations in both pasture yield and composition, particularly of annual and ephemeral species growing between these perennial tussocks (Orr and Holmes 1984). These fluctuations create problems when attempting to assess rangeland condition and trend which is often based upon single, or short term, data. Furthermore, Everist and Webb (1975) concluded "the extrapolation from observations made at any one time can be misleading and inaccurate". This paper summarises an approach by which State and Transition theory (Westoby *et al.* 1989) and the Degradation Gradient Method (Bosch and Kellner 1991) were used to develop models of vegetation change which were able to account for this inherent variability.

VEGETATION SURVEYS

One hundred and sixty two survey sites, 30 by 30 m square were permanently marked to represent different conditional states (fence line effects, piosphere transects and stock camps) within relatively homogeneous grazing areas (RHGAs) of the Mitchell grasslands in Queensland. Sites were surveyed in August 1989, January 1990, May 1990, May 1991, May 1993 and May 1994. The nearest plant within 200 mm of a set strike was recorded approximately every 1 m. Additional data, such as soil pH, topography and erosion activity were determined for each site at the commencement of the study with pasture biomass and grazing pressure estimated on each survey date (Phelps 1999).

DEFINING STATES, TRANSITIONS AND DEGRADATION GRADIENTS

Vegetation data from each recording date was ordinated separately in a time-series approach (ie each survey date analysed individually) to avoid rainfall patterns unduly influencing the proportional plant abundance data. The ordination techniques, Decorana and Reciprocal Averaging, were used to identify different vegetation groups (Bosch *et al.* 1992). These groups were treated as states within each recording date. The movement of each site within a state was traced over time. Most sites retained their grouping from date to date, providing evidence of generalised vegetation states. Some sites moved from one state to another over time, providing evidence of transitions. These states and transitions were then combined to form a generalised State and Transition model for each RHGA.

The existence of a degradation gradient was then confirmed or rejected by plotting edaphic factors and known management histories (eg protected sites, sites with known utilisation levels within grazing trials, sites near watering points and stock camps) on the ordination diagrams of each date. This defined the main ecological factor(s) responsible for the positioning of the states along the first axis. For example, a degradation gradient was defined for the Longreach district (Figure 1). Vegetation states ranged from the perennial grassland of State I (good condition) to the less stable herbfield states dominated by forbs and bare ground (poor condition). This was confirmed by the grazing histories of a number of sites within the ordinations, the extent of surface litter, erosion activity, and bare ground. Most lightly utilised sites occurred in State I, while the light to moderate and moderate to highly utilised sites occurred in State II. Sites that were overgrazed and/or severely disturbed (near watering points, stock camps and in overgrazed corners) occurred only in the herbfield states. Erosion activity was also highest in the herbfield states, while the amounts of litter were highest in States I and II.




Figure 1. A simplified State and Transition model for the Mitchell grasslands of the Longreach district (from Phelps 1999).

DETERMINING INDICATOR SPECIES

If a degradation gradient existed, then indicator species could be identified. Firstly, species abundances were plotted along the time specific vegetation states and Gaussian fit regression analyses performed. Where rainfall patterns were similar at different dates (eg rains following a seasonal drought) similarities were also found in both species abundance and behaviour. This made it possible to summarise the indicator species according to rainfall patterns. Key indicator species of the states along the gradient were identified as those with high r^2 values, and with relatively high abundances.

CONCLUSION

This approach was effective in quantitatively defining states, transitions, degradation gradients and indicator species for the Mitchell grasslands. These models can now be used to interpret data from other vegetation monitoring programs. Our model further develops the general State and Transition model of McArthur *et al.* (1994) through the use of quantitative data collected in smaller areas of Mitchell grassland.

REFERENCES

- Bosch, O.J.H. and Kellner, K. (1991). The use of degradation gradients for the ecological interpretation of range condition assessment. *J. Arid. Environ.* 21: 21-29.
- Bosch, O.J.H., Gauch, H.G., Booyesen, J., Stols, S.H.E., Gouws, G.A., Nel, M.E. and Van Zyl, E. (1992). 'Integrated System for Plant Dynamics', Ver 1.0. Copyright (C) 1991, Dept. Plant and Soil Sciences, P.U. for CHE, Potchefstroom, South Africa.
- Everist, S.L. and Webb, L.J. (1975). 'Two communities of urgent concern in Queensland: Mitchell grass and tropical closed forests. 2. National system of ecological reserves in Australia'. *Aust. Acad. Sci. Rep. No. 19*: pp 39-52.
- McArthur, S R. Chamberlain, H J. and Phelps, D G. (1994). A general state and transition model for the Mitchell grass, bluegrass/browntop and Queensland bluegrass pasture zones of northern Australia. *Trop. Grassl.* 28: 274-8.
- Orr, D.M. and Holmes, W.E. (1984). Mitchell grasslands. In 'Management of Australia's Rangelands' (Eds G.N. Harrington, A.D. Wilson and M.D. Young). CSIRO Melbourne pp 241-254.
- Phelps, D.G. (1999). 'Monitoring the Rangelands of Western Queensland'. Department of Primary Industries, Queensland. QO99019.
- Westoby, M., Walker, B. and Noy-Meir, I. (1989). Opportunistic management for rangelands not at equilibrium. *J. Rangel. Manage.* 42: 266-274.