

PROCEEDINGS OF THE AUSTRALIAN RANGELAND SOCIETY BIENNIAL CONFERENCE
Official publication of The Australian Rangeland Society

Copyright and Photocopying

© The Australian Rangeland Society. All rights reserved.

For non-personal use, no part of this item may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior permission of the Australian Rangeland Society and of the author (or the organisation they work or have worked for). Permission of the Australian Rangeland Society for photocopying of articles for non-personal use may be obtained from the Secretary who can be contacted at the email address, rangelands.exec@gmail.com

For personal use, temporary copies necessary to browse this site on screen may be made and a single copy of an article may be downloaded or printed for research or personal use, but no changes are to be made to any of the material. This copyright notice is not to be removed from the front of the article.

All efforts have been made by the Australian Rangeland Society to contact the authors. If you believe your copyright has been breached please notify us immediately and we will remove the offending material from our website.

Form of Reference

The reference for this article should be in this general form;

Author family name, initials (year). Title. *In*: Proceedings of the *n*th Australian Rangeland Society Biennial Conference. Pages. (Australian Rangeland Society: Australia).

For example:

Anderson, L., van Klinken, R. D., and Shepherd, D. (2008). Aerially surveying Mesquite (*Prosopis* spp.) in the Pilbara. *In*: 'A Climate of Change in the Rangelands. Proceedings of the 15th Australian Rangeland Society Biennial Conference'. (Ed. D. Orr) 4 pages. (Australian Rangeland Society: Australia).

Disclaimer

The Australian Rangeland Society and Editors cannot be held responsible for errors or any consequences arising from the use of information obtained in this article or in the Proceedings of the Australian Rangeland Society Biennial Conferences. The views and opinions expressed do not necessarily reflect those of the Australian Rangeland Society and Editors, neither does the publication of advertisements constitute any endorsement by the Australian Rangeland Society and Editors of the products advertised.



The Australian Rangeland Society

The risky business of designing land tenure policy in the Gobi Desert

Addison, J.^{1,2}, Friedel, M.¹, Davies, J.¹, Tiver, F.³ and Bastin, G.¹

¹CSIRO Sustainable Ecosystems, PO Box 2111, Alice Springs, NT, 0871, jane.addison@csiro.au

²University of Queensland, School of Integrative Systems, St Lucia, QLD, 4067

³University of South Australia, Mawson Lakes, South Australia, 5095

Keywords: variability; pastoral; climate

Abstract

Pastoralists need to manage the feed gap risk that climatic/forage resource variability presents. We use the PHYGROW model to better understand interactions between this variability and the effectiveness of a geographically fixed pastoral community management area (CMA) in the Gobi Desert. The model shows significant intra- and inter-annual forage variability within sites, as well as significant variation between sites. Interviews with 9-Erdene members, and others, indicate that 'rule breaking' through departure of pastoralists and livestock from the area during frequent poor forage years is common, although other CMA institutions that maximise livestock body condition have been maintained. There is significant risk that designing and adequately policing geographically fixed pastoral systems at this scale will facilitate overgrazing and reduce livelihoods. The provision of financial and in-kind support that provides more diverse, affordable and accessible risk management options to pastoralists may be more effective at promoting good rangeland condition than further effort to closely define the boundaries of household grazing lands.

Introduction

Land tenure policies have been recognised as a significant driver of pastoralist decision-making in arid rangelands e.g. Productivity Commission (2002). Pastoralists need to respond to the feed gap risk that extreme climatic and resource variability presents within the constraints imposed by prevailing land use institutions. Here the Mongolian Gobi Desert's

9-Erdene community management area (CMA), which has a geographically defined boundary, is used to demonstrate how significant spatio-temporal forage variability can interact with areal restrictions to increase the risk of pastoralists experiencing significant forage feed gaps. This paper gives interim results of PhD research, and as such offers a ‘snap-shot’ only.

Materials and methods

The 9-Erdene CMA (Fig. 1) was selected in 2009 as interviews with NGO officials suggested that the group was functional and well-established, thus likely to have experienced more climatic variation than other such areas. Whilst more detailed socioeconomic surveys will be made during the 2010 field season, semi-structured, informal interviews with pastoralists, local officials, non-government organisations and researchers were undertaken between 2007 and 2009. These interviews elicited CMA geographical size, age and establishment information, formal and informal institutions governing pastoralist movements and the type and level of ‘rule breaking’ of such institutions.




Fig. 1. 9-Erdene CMA location, including the four sites where the PHYGROW model was verified. The photo is of verification site UG9E01, June 2009.

PHYGROW (Texas A & M University), a forage model linked to real-time climate data, was utilised to better understand spatial and temporal variations in forage availability within

9-Erdene. Four sites were selected within the CMA to be at least 5km apart and at different altitudes to maximise variability captured by the model. Aspect, slope, altitude, % bare ground, % litter, % vegetation cover, and the non-woody clipped, oven-dried and weighed vegetation of ten 0.5m² quadrats per site were verified in the field as per Rowan (1995). This field data complemented a combination of remotely sensed data, the literature, local and international expert knowledge and pre-existing soil maps (source: Institute of Geoecology, Mongolia) to inform other parameters.

Results

The 9-Erdene CMA, established in 2002 by the Initiative for People Centred Conservation of the New Zealand Nature Institute, covers 367km² (NZNI, 2006). Most pastoralists have rights over winter/spring shelters that are recognised legally. It is unclear whether winter/spring shelters are transferrable either under national law or group conventions, although evidence from similar CMAs indicate pastoralists are increasingly gaining rights over multiple shelters to sub-lease during times of limited forage elsewhere (Addison, unpublished). Similarly, whilst exclusivity is implied with group membership, exclusivity under national law is unclear and group boundaries are porous in reality. Pastoralists in this area currently manage climatic risk through mobility, alternative incomes (Upton 2003) and livestock banks.

Modelled forage availability shows significant intra- and inter-annual variability (Fig. 2). A late summer/autumn peak is consistent each year but the magnitude of the peak varies significantly. Despite significant inter-annual variability in precipitation over winter/spring, forage is much less variable during this period because temperatures are too low to support plant growth, irrespective of precipitation levels.




Fig. 2. Simulated total standing crop (kg/ha) at four sites, with altitudes, within 9-Erdene. 'Buffering risk' is further explained in the text.

'Buffering risk' (Fig. 2) is a comparative estimate of the maximum potential forage available to pastoralists if they can freely move within 9-Erdene, with interviews suggesting pastoralists largely follow this pattern. In winter, pastoralists generally stay in camps protected from wind e.g. UG9E01, moving in summer to areas with water and cooling winds e.g. UG9E04. Pastoralists return to the same winter camp in most years, but show much more variability in camp selection in summer, both within and between years. In low forage years (like 2009) pastoralists also move out of the CMA altogether, breaking group agreements. Pastoralists also cite receiving other non-member pastoralists from outside the geographically constrained 9-Erdene area when there is a lack of forage in surrounding areas, such as in the 'dzud' years of 2000/2001 (Upton, 2003). By following this pattern, the more rainfall-dependent, short-term vegetation pulses of the plains are utilised in the more variable summer, resting the more perennial and shrub-like forage utilised over winter. Available forage is therefore greater following the 'buffering risk' pattern (with occasional migrations outside the 9-Erdene area) than if a pastoralist was to stay all-year at any one site, other than the high altitude UG9E01. Staying full-time at high altitude sites is likely to degrade the pastoral resource held in common, however, so group members have maintained the CMA practise of an annual meeting to set group departure dates from winter/spring camps.

Discussion

Whilst Ostrom (2008) warns that there are no 'optimal' rules applicable to common pool resources, she suggests that CMAs can be effective if they are tailored to specific resource attributes. Many of the critical enabling conditions for sustainability of the commons, synthesised by Agrawal (2003), are not met in the non-equilibrium 9-Erdene CMA. However, the forage resource is neither predictable nor immobile, nor does it have a well-defined boundary. Given the frequent 'rule breaking' of the CMA boundary during low forage years, it is likely that there has been a mismatch between the spatio-temporal scale of the forage resource and the areal extent of the CMA. That 9-Erdene is one of the few functional CMAs left in the area suggests that the group has gained value from being selective in the aspects of the formalised system it has adopted, retaining specific aspects of the CMA, like spring/winter leaving dates, whilst dropping unhelpful elements, like fixed boundaries. It is probable that the latter do not support the aim of fattening livestock over summer/autumn to help withstand the long winter.

The lag-time between rainfall-mediated vegetation pulses and the build-up of significant grazing pressure in non-equilibrium rangelands is greater in mobile systems, and/or where livestock numbers are not maintained through artificial waterpoints or supplementary feeding during poor rainfall seasons (Vetter, 2005). This lag-time is nearly non-existent in the geographically constrained 9-Erdene. The adequately policed restricting of pastoralists to such areas therefore increases the risk of overgrazing, particularly of high value, summer-available species like *Allium sp* and, ultimately, a decline in livelihoods. These are key concerns of both governments and multilateral organisations that, understandably, face difficulties in designing suitable programmes and policies for the region. However, the provision of financial and in-kind support that provide mores diverse, affordable and accessible risk management options to pastoralists are likely to be more effective in the short to medium term than further effort to enforce the temporal and spatial boundaries of household grazing lands.

Acknowledgements

The authors are very grateful to the Australian Rangeland Society for partially funding the 2009 field season. The Desert Knowledge CRC and Endeavour Research Fellowship also

provided financial support for this work. MercyCorps Mongolia and Texas A & M University provided other valuable assistance.

References

Agrawal, A. (2003) Sustainable governance of common-pool resources: context, methods and politics. *Annual Review of Anthropology* **32**, 243-262.

MercyCorps Mongolia. (2004) Gobi Forage Project Information. Available online: www.mercycorps.org.mn/index.

New Zealand Nature Institute. (2006) Summary of Project Support Strategies and Impacts on Local Livelihoods and Natural Resources. Draft Report. January 2006.

Ostrom, E. (2008) Institutions and the environment. *Economic Affairs* **28**, 24-31.

Productivity Commission (2002) 'Pastoral leases and non-pastoral land use.' (Commission Research Paper, AusInfo: Canberra).

Rowan, R. (1995) PHYGROW model documentation version 2. Rancing System Group, Dept Rangeland Ecology Management, Texas A & M University, College Station.

Upton, C. (2003) Tragedy or Transformation? Institutions, Collective Action and Conservation on Mongolia's Pastoral Commons. Ph.D Thesis, University of Cambridge, Cambridge, England.

Vetter, S. (2005) Rangelands at equilibrium and nonequilibrium: recent developments in the debate. *Journal of Arid Environments*. **62**, 321-341.

Addison, J., Friedel, M., Davies, J., Tiver, F., Bastin, G. (2010). The risky business of designing land tenure policy in the Gobi Desert. In: *Proceedings of the 16th Biennial Conference of the Australian Rangeland Society*, Bourke (Eds D.J. Eldridge and C. Waters) (Australian Rangeland Society: Perth).