

PROCEEDINGS OF THE AUSTRALIAN RANGELAND SOCIETY BIENNIAL CONFERENCE
Official publication of The Australian Rangeland Society

Copyright and Photocopying

© The Australian Rangeland Society. All rights reserved.

For non-personal use, no part of this item may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior permission of the Australian Rangeland Society and of the author (or the organisation they work or have worked for). Permission of the Australian Rangeland Society for photocopying of articles for non-personal use may be obtained from the Secretary who can be contacted at the email address, rangelands.exec@gmail.com

For personal use, temporary copies necessary to browse this site on screen may be made and a single copy of an article may be downloaded or printed for research or personal use, but no changes are to be made to any of the material. This copyright notice is not to be removed from the front of the article.

All efforts have been made by the Australian Rangeland Society to contact the authors. If you believe your copyright has been breached please notify us immediately and we will remove the offending material from our website.

Form of Reference

The reference for this article should be in this general form;

Author family name, initials (year). Title. *In*: Proceedings of the *n*th Australian Rangeland Society Biennial Conference. Pages. (Australian Rangeland Society: Australia).

For example:

Anderson, L., van Klinken, R. D., and Shepherd, D. (2008). Aerially surveying Mesquite (*Prosopis* spp.) in the Pilbara. *In*: 'A Climate of Change in the Rangelands. Proceedings of the 15th Australian Rangeland Society Biennial Conference'. (Ed. D. Orr) 4 pages. (Australian Rangeland Society: Australia).

Disclaimer

The Australian Rangeland Society and Editors cannot be held responsible for errors or any consequences arising from the use of information obtained in this article or in the Proceedings of the Australian Rangeland Society Biennial Conferences. The views and opinions expressed do not necessarily reflect those of the Australian Rangeland Society and Editors, neither does the publication of advertisements constitute any endorsement by the Australian Rangeland Society and Editors of the products advertised.



The Australian Rangeland Society

Biodiversity impacts of ground tank closure in southeast Australian rangelands

Westbrooke, M.E. ¹, Christie, F.J. ^{1,2}, Cook, S. ¹, Florentine, S.K. ¹, Gell, P. ¹ and Graz, P. ¹

¹Centre for Environmental Management, University of Ballarat

²Centre for Ecosystem Science, University of Melbourne (current)

Keywords: Biodiversity; water point; grazing pressure

Abstract

Literature suggests that provision of artificial water through bores, dams and ground tanks has negative environmental effects through increased grazing pressure, prevention of native species regeneration and promotion of exotic species. Though some native species may benefit from increased water, other species may decline. In arid environments worldwide, few areas remain distant from water. In Australia, much water provision is through ground tanks storing diverted run-off following rainfall events. Whilst closure of tanks is a recognized means of reducing negative impacts there is reluctance to utilize this strategy. Consequently, there are few opportunities to study responses of plant and animal populations to such closures. Where tank closure has occurred in areas reserved for conservation there is a lack of data on the effects on flora and fauna. Past studies have been limited in: the range of biota incorporated; lack of landscape-scale experimental manipulation; failure to replicate treatments. Research on the University of Ballarat's 40,000 hectare arid-zone research property, Nanya, is investigating impacts of four alternative tank treatments: fencing to exclude mammalian grazers; partial tank closure by blocking of drains; complete closure with landscaping to original profile; no action. Responses of plants, mammals, birds, reptiles and ants to these treatments are being studied. Findings will help illuminate the potential for water point closure to restore balance to the surrounding landscape, an important consideration in biodiversity conservation in arid landscapes in Australia and other parts of the world

Introduction

Water is a key limiting factor in arid environments and, prior to European settlement Australian arid zones had few permanent water points (James *et al.* 1995). Abundant literature suggests that provision of artificial water through bores, dams and ground tanks in pastoral areas has had negative environmental effects (e.g. Landsberg *et al.* 1997; Hilty *et al.* 2006, Todd 2006). Currently there are few areas left in the Australian rangelands that are distant from water, and this problem exists in arid and semi-arid environments worldwide (e.g. Noy-Meir 1996, Thrash 1998).

Some of the potential impacts of watering points on the surrounding environment include:

- Contributing to elevated native and exotic grazer populations (James *et al.* 1995)
- Redistribution of nutrients through deposition of dung (Lange & Willcocks 1978)
- Allowing establishment and spread of exotic plant species (Westbrooke 1990) and unpalatable native plant species.
- Prevention of regeneration of native perennial plant species (Westbrooke 2005), in particular palatable chenopods (Montague-Drake & Croft 2004)
- Increased numbers of opportunistic water-dependent species (Draper *et al.* 2004).
- Providing a focus for activity of exotic predators
- Potential flow on effects for tree health through changes in trophic interactions (Landsberg *et al.* 1990, Grey *et al.* 1997).

Fensham and Fairfax (2008) in reviewing this issue highlighted the lack of sound data on benefits of closing water-points and the need to address the problem at a landscape scale. In western NSW and adjacent areas of South Australia, a major means of water provision is through ground tanks storing diverted run-off following major rainfall events. There is widespread agreement that, for better control of feral grazers and improved grazing management in areas where biodiversity is a primary or secondary objective, closure of tanks or greater control of water distribution is management strategy. Despite incentives available for improving biodiversity outcomes on pastoral lands, there is reluctance among landholders to utilise these strategies and as a result, there are few opportunities to study in detail the responses of plant and animal populations to such closure. Past studies have not

carried out experimental manipulation at the landscape scale to demonstrate the long-term impacts on arid-zone biodiversity.

Research is investigating the impacts of four alternative tank treatments on a range of plant and animal groups.

Methods

The research is centered on Nanya Station (40,000 ha.) 150 km northwest of Wentworth in southwestern NSW (Fig. 1), where the dominant vegetation comprises eucalypt open-scrub (mallee) (Westbrooke *et al.* 1998). Until 2004, Nanya Station was under pastoral management but its purchase by the University of Ballarat for teaching, research and conservation provides a unique opportunity to test the conservation benefits of manipulating water availability across the landscape.




Fig. 1. Location of study area * indicates Nanya Station

Twelve tanks ground tanks were selected for maximum distance between treatments, similarity in size of tank and surrounding vegetation type. Assessments are undertaken at sites adjacent and at 500m and 1500m from the tank, replicated on dunes and in swales (Fig. 2).




Fig. 2. Simplified diagram indicating experimental design and location of assessment sites at each of 12 ground tanks

At all sites, a range of taxa was surveyed prior the application of treatments (fencing to exclude mammalian grazers; partial tank closure by blocking of drains; complete closure with landscaping to original profile; no action – each with three replicate tanks) and at intervals following treatment. These surveys include:

Large grazing mammals: Assessed annually through faecal accumulation plots. Plots are checked for faeces accumulated during each assessment, and the number of faecal groups for each species identified, following the technique of Neff (1968).

Reptiles and small mammals: Abundance and richness assessed in spring, summer and autumn via 20-meter pitfall lines. Pitfall lines are run for five nights during each assessment.

Predators: Fox scats are collected and analysed for hair and bone content and level of fox activity is assessed based on uptake of baits which are set on a 1km. grid at six monthly intervals.

Birds: Species abundance and richness assessed in spring and autumn through morning 20min. point survey counts.

Ants: Surveyed via pitfall trapping in summer to assess changes in species composition of the community.

Vegetation: The line intercept of all ground species along with litter, cryptogamic crust and bare ground is sampled along transect lines within each quadrant. The point-centred quarter

technique is used to determine the occurrence and density of tree and shrub species.

Structural attributes are assessed for a 50 x 20m area around the pitfall line.

Climate: Since rainfall in the area is highly variable both temporarily and spatially rainfall and soil moisture are being assessed by automatic weather stations established at each water point.

Results

For the vegetation, preliminary results indicate a positive relationship (cover and richness) between native species and distance from water and a negative relationship for exotic species. These trends were more marked for dunes than swales.

There is higher large mammal activity at water points but whilst goats are more abundant in swales, Grey Kangaroos prefer dunes whereas Red Kangaroos show no habitat effect.

A total of 93 bird species have been recorded during surveys. Two of these were EPBC listed species and a further four listed under state legislation. Greater species richness was detected within 20m of water however there were significant compositional differences with more parrots and honeyeaters but less small insectivores. Additionally, there was a significant increase in the total abundance of ground nesting birds (Southern Scrub Robin, Malleefowl, Chestnut Quail Thrush and Striated Grass Wren) recorded with increasing distance from water.

Forty-seven species of reptile and six species of small mammals were collected via pitfalls. Water points have less reptile species than other sites. Dunes have greater abundance than swales and tanks but all three habitats have a distinct complement of species. Sites 500m< from water had a greater abundance of small mammals than sites adjacent to watering points. Preliminary analysis indicates that ant diversity may be greater in close proximity to tanks than at other sites.

Discussion

Limited time since treatment and variability of rainfall make it too early to draw conclusions on the relative impact of the treatments. However 18 months post treatment) and based on pre treatment surveys, taxa have shown highly variable responses to distance from water. Ground nesting bird species are likely to be exposed to risks from predation, particularly in more open landscapes such as those which exist in close proximity to watering points. This pattern was reflected in our data and is closely related to a reduction in structural complexity of vegetation and habitat closer to watering points as demonstrated through vegetation surveys and habitat assessments. Preliminary vegetation data support the view that, in conservation reserves created in areas previously used for pastoralism, an essential measure for vegetation recovery is closure of watering points.

References

Draper, R., Gowans, S., Gibson, M., Baker, B. and Milne, R. (2004). Wetland Dependant Flora in the Birchip Area, North West Victoria. CHEC and CEM, University of Ballarat.

Fensham, R.J. and Fairfax, R.J. (2008). Water-remoteness for grazing relief in Australian arid-lands. *Biological Conservation* **141**, 1447-1460.

Grey M.J., Clarke M.F. and Loyn R.H. (1997). Initial changes in the avian communities of remnant eucalypt woodlands following a reduction in the abundance of noisy miners, *Manorina melanocephala*. *Wildlife Research* **24**, 631-648.

Hilty, J. A., Brooks, C., Heaton, E., and Merenlender, A.M. (2006). Forecasting the effect of land-use change on native and non-native mammalian predator distributions. *Biodiversity and Conservation* **15**, 2853-2871.

James, C.D., Landsberg, J. and Morton, S.R. (1995). Ecological functioning in arid Australia and research to assist conservation of biodiversity. *Pacific Conservation Biology* **2**, 126-142.

Landsberg, J, James, C., Morton, S., Hobbs, T., Stol, Drew, A. and Tongway, H. (1997). The Effects of Artificial Sources of Water on Rangeland Biodiversity. Final report to the Biodiversity Convention and Strategy Section of Environment Australia, Canberra.

Lange, R. and Willcocks, M.C. (1978). The relation between sheep-time spent and egesta accumulated within an arid-zone paddock. *Australian Journal of Experimental Agriculture and Animal Husbandry* **18**, 764-767.

Montague-Drake, R. and Croft, D.B., 2004. Do kangaroos exhibit waterfocussed grazing patterns in arid New South Wales? *Australian Mammology* **26**, 87–100.

Neff, D.J. (1968). The pellet-group count technique for big game trend, census, and distribution: a review. *Journal of Wildlife Management*. **32**, 597- 614

Noy-Meir, I, (1996) The spatial dimensions in plant –herbivore interactions. In: West, N.E. (Ed.) Rangelands in a sustainable Biosphere. *Proceedings of the Fifth International Rangeland Congress*, Volume 2, pp 152-154. Denver, CO: Society for Range Management.

Thrash, I. (1998) Impact of water provision on herbaceous vegetation in Kruger National Park, South Africa. *Journal of Arid Environments*, **38**, 437-450.

Todd, S. W. (2006). Gradients in vegetation cover, structure and species richness of Nama-Karoo shrublands in relation to distance from livestock watering points. *Journal of Applied Ecology* **43**, 293-304.

Westbrooke, M.E. (1990). The effects of grazing on weediness in Mallee communities - studies at Mallee Cliffs National Park and Nanya Station, Western N.S.W. In J. Noble, eds. *The Mallee Lands - a Conservation Perspective*, CSIRO: Melbourne.

Westbrooke, M.E. (2005). Relationship between perennial species richness and distance from water in *Casuarina pauper* (Belah) woodland. *Victorian Naturalist* **122**, 57-62.

Westbrooke, M.E., Miller, J.D. and Kerr, M.K. (1998). Vegetation and flora of the Scotia 1:100 000 map sheet, far western New South Wales. *Cunninghamia* **5**, 665-684.

Westbrooke, M.E., Christie, F.J., Cook, S., Florentine, S.K., Gell, P. and Graz, P. (2010). Biodiversity impacts of ground tank closure in southeast Australian rangelands. In: *Proceedings of the 16th Biennial Conference of the Australian Rangeland Society*, Bourke (Eds D.J. Eldridge and C. Waters) (Australian Rangeland Society: Perth).