

PROCEEDINGS OF THE AUSTRALIAN RANGELAND SOCIETY BIENNIAL CONFERENCE
Official publication of The Australian Rangeland Society

Copyright and Photocopying

© The Australian Rangeland Society. All rights reserved.

For non-personal use, no part of this item may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior permission of the Australian Rangeland Society and of the author (or the organisation they work or have worked for). Permission of the Australian Rangeland Society for photocopying of articles for non-personal use may be obtained from the Secretary who can be contacted at the email address, rangelands.exec@gmail.com

For personal use, temporary copies necessary to browse this site on screen may be made and a single copy of an article may be downloaded or printed for research or personal use, but no changes are to be made to any of the material. This copyright notice is not to be removed from the front of the article.

All efforts have been made by the Australian Rangeland Society to contact the authors. If you believe your copyright has been breached please notify us immediately and we will remove the offending material from our website.

Form of Reference

The reference for this article should be in this general form;

Author family name, initials (year). Title. *In*: Proceedings of the *n*th Australian Rangeland Society Biennial Conference. Pages. (Australian Rangeland Society: Australia).

For example:

Anderson, L., van Klinken, R. D., and Shepherd, D. (2008). Aerially surveying Mesquite (*Prosopis* spp.) in the Pilbara. *In*: 'A Climate of Change in the Rangelands. Proceedings of the 15th Australian Rangeland Society Biennial Conference'. (Ed. D. Orr) 4 pages. (Australian Rangeland Society: Australia).

Disclaimer

The Australian Rangeland Society and Editors cannot be held responsible for errors or any consequences arising from the use of information obtained in this article or in the Proceedings of the Australian Rangeland Society Biennial Conferences. The views and opinions expressed do not necessarily reflect those of the Australian Rangeland Society and Editors, neither does the publication of advertisements constitute any endorsement by the Australian Rangeland Society and Editors of the products advertised.



The Australian Rangeland Society

WATERSPREADING TO RESTORE NATIVE GRASSLANDS

Kevin Mitchell¹

¹Property Landholder
“Florida” Canbelego New South Wales 2835 Australia
Email: ray.thompson@cma.nsw.gov.au

Matthew Tighe²

²Research Fellow Ecosystem Management
University of New England, Armidale NSW 2351
Email: mtighe2@une.edu.au

Ray Thompson³

³Central West Catchment Management Authority
PO Box 121 Nyngan New South Wales 2825 Australia
Email: ray.thompson@cma.nsw.gov.au

“Key words:” Waterspreading. Environmental. Invasive native scrub. Land management. Soil carbon.

Abstract

Waterspreading is a land rehabilitation technique that targets the variability of rainfall and runoff in semi-arid systems to initiate long term changes in ground cover. This study outlines the effect of waterspreading at ‘Florida’ in western NSW, which has been steadily implementing waterspreading systems for the last 30 years. By combining recent pasture measurements, on-farm observations, and soil surface carbon and nitrogen measurements, this study outlines the dramatic changes in pasture condition and diversity, and long term changes in surface soil properties, that occur following waterspreading. These dramatic yet persistent changes exemplify the benefits of implementing rehabilitation that is based upon the processes that govern resource movement and productivity within semi-arid systems, namely, recognition of variability in rainfall and runoff, and management of this.

Introduction

Before European settlement, the Western New South Wales Peneplain landscape was a mosaic of open grasslands with patches of scrub and open woodlands – now Invasive Native Scrub dominates the landscape which has led to reduced carry capacity (Gardiner *et al.* 1998), such as reduced biodiversity and changes in soil condition (Tighe *et al.* 2009). Many methods have been tried to bring these degraded areas of the peneplain back to production. Waterspreading is one method that is very successful.

Waterspreading is a land management technique used to evenly spread and disperse rainwater flows over country with gentle slopes less than three percent. It is based on the recognition of the patchiness of semi-arid landscapes, and the source-sink operation of different soil surface conditions (Tongway and Ludwig 1996). The driving mechanism behind the success of waterspreading is the reduction of the energy of water flow, meaning a large reduction in soil erosion and an increase in water infiltration.

Waterspreading involves creating a series of small banks to direct water away from eroding drainage lines to areas where it would normally not flow (Central West & Western CMA 2008). Each bank is designed to slow and spread water as it continues downslope, increasing infiltration as to better suit native grasses and herbage. Restoring native grassland from an INS site in this way is a long-term exercise and can take five to seven years, depending on the season and soil of the area (Central West & Western CMA 2008). Firstly the invasive scrub area is thinned or cleared through chaining, raking and burning, followed by ploughing if necessary because of regeneration of turpentine and bumble box. Waterspreading banks are surveyed and constructed. Pioneer plants such as yellow burr daisy and galvanised burr will establish first. These pioneer plants provide ground cover and

will be replaced by native grasses and herbage after around two years. While the changes in ground cover and herbage are evident after this time period, there has been little documentation of this. In addition, improvements in soil condition following waterspreading, there is noticeable change in colour as biomass increases. This paper presents a summary of initial visual responses of a grazing system to waterspreading, and surface soil measurements within this system that is indicative of soil condition.

Methods

Study site, survey and construction

Measurements and observations were undertaken on 'Florida', Canbelego, NSW, approximately 50 km east of Cobar. Over the last 30 years 500 kilometres of waterspreading banks have been constructed covering 3400 hectares on Florida. The general method of waterspreading construction is as follows: the paddock is surveyed. Following marking, roadgraders are used to construct waterspreading banks – preferably to a height of 60cm and basal width of over 2.0 metres – which are then rolled to account for stock trampling, tunnelling, flooding and settlement of bank.

Monitoring

Herbage mass and ground cover were documented in 2008 on an area untreated and treated north of the Florida homestead by Higgins and Theakston (2008). Changes in species diversity and composition and stocking rates are a summary of visual observations since the beginning of waterspreading trials on this property over 30 years ago (K. Mitchell, pers. obs.).

For soil analysis, three waterspreading systems were selected on 'Florida', 2, 15 and 30 years since waterspreading implementation, and matching site conditions as closely as possible. An adjacent patch of scrub (dominated by Turpentine – *Eremophila sturtii*) taken to be representative of pre-spreading conditions was also sampled. Surface (0-5cm) soil sampling for carbon (C), nitrogen (N), and available N (the sum of NH_4^+ N and NO_3^- N) and statistical analysis followed the approach of Tighe *et al.* (2009), with Tukey's posthoc comparisons undertaken where significant differences ($P < 0.05$) were detected.

Results

Pasture dynamics

Table 1. Average herbage dry weight and ground cover at adjacent untreated site and after waterspreading. Samples taken in 2008 across an area that was Chained 2002, Raked & Waterspread 2006, Two years under a waterspreading systems.

Variable	Before Waterspreading	After Waterspreading
Herbage dry weight (units)Kg/Ha	146	5,061
Ground cover (%)	26	98
Visual change in pasture species across whole waterspreading system	80% No9 Wiregrass & Corkscrew	5% No9 Wiregrass & Corkscrew
Soft palatable grasses & herbage	5%	80%
Estimated sustainable stocking rate	1 cow to 50 hectares	1 cow to 9 hectares
Species diversity	47	+ 350

Table 1 shows a pronounced increase in both average herbage dry weight and ground cover following water spreading. Similarly, there has been a very obvious shift away from the dominance of undesirable pasture species, and an increase in pasture species diversity. These changes have resulted in a notable improvement in the on-farm stocking rate.

Soil C and N

Total C, and available N showed similar trends across the different ages of waterspreading systems sampled, compared with the scrub patch (Fig. 1). Total N followed C very closely and is not reported here. The 2 year old waterspreader had higher C, N, and available N compared with the scrub patch. This difference was not evident in the 15 year old system, but was present in the oldest (30 year old) system. In the 30 year old system the available N was significantly higher compared to all other systems.




Fig. 1. Total carbon (A) and available N (B) in 0-5 cm soil of three waterspreading systems of different ages, and one adjacent patch of Turpentine (*E. sturtii*) mixed scrub. Different lowercase letter denote differences between sites at $P < 0.05$.

Discussion

There are complex processes operating in semi-arid systems, processes that waterspreading capitalises on. By building upon an understanding of how runoff interacts with soil condition, waterspreading targets the source-sink functioning of the soil surface over a timeframe that is relevant to the boom-bust cycles of the semi-arid zone. The difference between waterspreading treated and non-treated INS sites is clear cut. The implementation of waterspreading banks combined with the removal of the Invasive Native Scrub (INS) is restoring grasslands on the landscape, improving pasture biodiversity, increasing carrying capacity and ensuring farm viability.

There is large potential for waterspreading to not only improve pasture stability and productivity, but to beneficially alter surface soil condition over a long time period. The higher C and available N in surface soils in the young water spreading system compared with the patch of scrub probably reflects the short term release of nutrients into the soil from biomass following the management of scrub (and burning of material) during waterspreader implementation. Similarly, the lower nutrient values in the 15 year old system may be indicative of this nutrient loss in the longer term, as the system utilises these new found, relatively available resources, or resources are removed via grazing or leaching. If this is the case, then the much higher values found in the 30 year old system may indicate the progression of the waterspreaders into a new stable state, in which nutrients accrue and cycle at higher levels. Similar results have been found in pastures of these regions following scrub management (Tighe *et al.* 2009), although the doubling of 0-5 cm C when the scrub patch is compared with the 30 year old waterspreader is a large comparative change compared with other grazing systems.

It is apparent from this study that waterspreading can produce dramatic changes in a degraded system, and it does so by targeting the processes that govern productivity and resource movement within semi-arid systems.

References

- Central West CMA & Western CMA, 2008. Waterspreading and restoring native grasslands on 'Florida'. Invasive Native Scrub Case Study: Waterspreading and INS management. Central West and Western Division Catchment Management Authorities, NSW.
- Gardiner, D.B., Tupper, G.J., Dudgeon, G.S., 1998. A quantitative appraisal of woody shrub encroachment in western New South Wales. *Rangeland J.* 20, 26-40.
- Higgins, C., Theakston, P., 2008. Paired Vegetation Comparisons Treated vs Untreated on "Florida Station" Western Catchment Management Authority, NSW.
- Tighe, M., Reid, N., Wilson, B.R., Briggs, S.V., 2009. Invasive native scrub and soil condition in semi-arid south-eastern Australia. *Agric. Ecosyst. Environ.* 132, 212-222.
- Tongway, D.J., Ludwig, J.A., 1996. Rehabilitation of semiarid landscapes in Australia. I. Restoring productive soil patches. *Restor. Ecol.* 4, 388-397.